

**▲ Lift Here**

For full prescribing information, see the product foldout.

Store at 59° - 86°F (15° - 30°C).

Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in the USP/NF with a child-resistant closure.

The labeling for this product includes professional labeling which is not intended for use by the general public.

**WARNING: KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.**

IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.



Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615

NDC 16477-401-01

**Cyclivert®**  
TABLETS

**Antiemetic**

For Healthcare Professionals:  
For the treatment of vertigo of motion sickness.

**Each tablet contains:**

**Cyclizine HCl** ..... 25 mg



**100 Tablets**

Rev 9/11



Lot No. :  
Exp. Date:

## Drug facts (continued)

Keep out of reach of children.

### Directions

Adults and children 12 years of age and over:	2 tablets every 4-6 hours, not to exceed 8 tablets in 24 hours.
Children 6 to under 12 years of age:	1 tablet every 6-8 hours, not to exceed 3 tablets in 24 hours.
Children under 6 years of age:	Consult a physician.

### Other information

Store at 59° - 86°F (15° - 30°C)

### Inactive ingredients

Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate

### Questions? Comments?

Rev 9/11

Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615



### Lift Here

For full prescribing information, see the product foldout.

Store at 59° - 86°F (15° - 30°C).

Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in the USP/NF with a child-resistant closure.

The labeling for this product includes professional labeling which is not intended for use by the general public.

**WARNING: KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.**

**IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.**

1.500

## Drug Facts

**Active ingredient Purpose**  
(in each tablet)  
Cyclizine Hydrochloride 25 mg  
..... Antiemetic

### Uses

For the prevention and treatment of the nausea, vomiting, or dizziness associated with motion sickness. For the treatment of vertigo of motion sickness.

### Warnings

**Do not exceed recommended dosage.**

**Ask a doctor or pharmacist before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor before use if you are** taking sedatives or tranquilizers.

**Do not give this product to children who have a breathing problem** such as chronic bronchitis or who have glaucoma, without first consulting the child's doctor.

**When using this product**

- you may get drowsy
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase the drowsiness effect
  - be careful when driving a motor vehicle or operating machinery
- If pregnant or breast-feeding, ask a health professional before use.**

**In case of overdose, get medical help or contact a Poison Control Center right away.**