

**FILE NAME: 1oz. Dallery Drops [ECL] Rev. 31.docx**

**NDC 16477-121-30**

**Dallery  
DROPS  
New Formulation**

**Sugar Free / Alcohol Free  
Antihistamine  
Nasal Decongestant**

**Peach Flavored**

Each dropperful (1.0 mL), for oral administration, contains:

**Chlorpheniramine Maleate ... 1.0 mg**

**Phenylephrine HCl ..... 2.5 mg**

**1 fl oz (30 mL)**

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Supplied in a tight, light-resistant container with a child-resistant closure.

Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615

Rev. 11/13

## Drug Facts

### **Active ingredients**

**(in each 1 mL dropperful)**

Chlorpheniramine Maleate, 1.0 mg

Phenylephrine Hydrochloride, 2.5 mg

### **Purpose**

Antihistamine

Nasal Decongestant

### **Uses**

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

### **Warnings**

**Do not exceed recommended dosage.**

#### **Do not use this product**

•if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### **Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

#### **When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- be careful when driving a motor vehicle or operating machinery

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **Do not exceed recommended dosage.**
- **Dosages may be repeated every 4 hours if required for relief.**
- **Use enclosed dropper only. Do not use any other dosage device.**

Adults and Children 12 years of age and over:	4 dropperfuls (4 mL) every 4 hours. Not to exceed more than 6 doses in 24 hours.
---	--

Children 6 to under 12 years of age: 24 hours.	2 dropperfuls (2 mL) every 4 hours. Not to exceed more than 6 doses in
--	---

Children under 6 years of age:	Consult a doctor.
-----------------------------------	-------------------

***Other information***

Store at 59° - 86° F (15° - 30° C)

***Inactive ingredients***

Citric Acid, FD&C Yellow #6, Glycerin, Peach Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

***Questions? Comments?***

Call your doctor for medical advice about side effects. Serious side effects associated with this product may be reported to this number. 1-864-286-8229 (Mon-Fri) 8am-5pm EST