

For full prescribing information, see the product foldout.

Store at 59° - 86°F (15° - 30°C)

Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in the USP/NF with a child-resistant closure.

The labeling for this product includes professional labeling which is not intended for use by the general public.

**WARNING: KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN. IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.**

NDC: 16477-820-01

# Dallergy<sup>®</sup>

## SYRUP

New Improved  
Formula

# LASER

Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615

Patent Pending

Rev. 08/2011



## Antihistamine Nasal Decongestant

Each 5 mL (1 teaspoonful) contains:

**Chlorcyclizine HCl ..... 12.5 mg**

**Phenylephrine HCl ..... 5 mg**



Grape Flavor

# LASER

**16 fl oz (473 mL)**

### **Drug Facts** (continued)

#### **Other information**

Store at 59° - 86°F (15° - 30°C)

#### **Inactive ingredients**

Bitter Mask, Citric Acid, FD&C Blue #1, FD&C Red #40, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sorbitol, Sucralose.

#### **Questions? Comments?**

Call 1-864-286-8229

# LASER

Manufactured for:  
Laser Pharmaceuticals, LLC  
Greenville, SC 29615

# Drug Facts

Lift Here

## Active ingredients Purpose (in each 5 mL teaspoonful)

Chlorcyclizine  
Hydrochloride 12.5 mg ..... Antihistamine  
Phenylephrine  
Hydrochloride 5 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing
- itching of the nose or throat
- itchy, watery eyes ■ nasal congestion
- reduces swelling of nasal passages

## Warnings

**Do not exceed recommended dosage.**

### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- heart disease ■ high blood pressure
- thyroid disease ■ diabetes

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers. ▶

4.125"

# Drug Facts (continued)

## When using this product

- excitability may occur, especially in children
- may cause drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- be careful when driving a motor vehicle or operating machinery

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

**Do not exceed 3 doses in a 24 hour period.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 6 to 8 hours, not to exceed 6 teaspoonfuls in 24 hours
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 6 to 8 hours, not to exceed 3 teaspoonfuls in 24 hours
Children 2 to under 6 years of age:	1/2 teaspoonful (2.5 mL) every 6 to 8 hours, not to exceed 1 1/2 teaspoonfuls in 24 hours
Children under 2 years of age:	Consult a doctor ▶