

FILE NAME: 100ct. Dallery Tablets Rev. 9.docx

NDC 16477-161-01

**Dallery
TABLETS
Improved Formulation**

Antihistamine • Nasal Decongestant

Each tablet contains:

Dexbrompheniramine Maleate 1 mg
Phenylephrine HCl 5 mg

100 Tablets

LASER

For full prescribing information, see the product foldout.
Store at 59° - 86°F (15° - 30°C).

Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing.

WARNING: KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.

IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

Manufactured for:
Laser Pharmaceuticals, LLC
Greenville, SC 29615

Manufactured by:
Pernix Manufacturing, LLC
Houston, TX 77099

Rev. 11/13

Drug Facts

Active ingredients (in each tablet)

Dexbrompheniramine Maleate
Antihistamine
Phenylephrine Hydrochloride
Nasal Decongestant

Purpose

1 mg
5 mg

Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

• runny nose • sneezing • itching of the nose or throat • itchy, watery eyes • nasal congestion • reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

• a breathing problem such as emphysema or chronic bronchitis • glaucoma
• difficulty in urination due to enlargement of the prostate gland • heart disease
• high blood pressure • thyroid disease • diabetes

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

• excitability may occur, especially in children • may cause drowsiness
• avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase the drowsiness effect • be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

• nervousness, dizziness, or sleeplessness occur • symptoms do not improve within 7 days or are accompanied by fever • new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children.

Directions

Do not exceed 6 doses in a 24 hour period.

Adults and children 12 years of age and over:	2 tablet every 4 hours, not to exceed 12 tablets in 24 hours.
---	---

Children 6 to under 12 years of age:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours.
---	--

Children under 6 years of age:	Consult a doctor.
-----------------------------------	----------------------

Other information

Store at 59° - 86°F (15° - 30°C)

Inactive ingredients

calcium phosphate dihydrate, colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, silicified microcrystalline cellulose

Questions? Comments?

Call 1-864-286-8229