

**Drug Facts** (continued)

**Directions**

- Do not exceed recommended dosage.
- Use only with enclosed dropper. Do not use with any other dosing device.

|                                      |   |
|--------------------------------------|---|
| Children 6 to under 12 years of age: | 2 dropperfuls (2 mL) every 4-6 hours, not to exceed 6 doses in 24 hours |
| Children under 6 years of age:       | Consult a Doctor  |

**Other information**

Store at 59° - 86°F (15° - 30°C)

**Drug Facts** (continued)

**Inactive ingredients**

Citric Acid, FD&C Red #40, Fruit Gum Flavor, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

**Questions? Comments?**

Call a doctor for medical advice about side effects. Serious side effects associated with use of this product may be reported to this number: 1-864-286-8229 (Mon-Fri) 8am-5pm EST

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.  
Supplied in a light, light-resistant container with a child-resistant closure.

Manufactured for:  
Esper Pharmaceuticals, LLC  
Grenada, SC 29616

Rev. 04/10

NDC 16477-811-30

**Donatussin  
DM DROPS**

**Antihistamine  
Antitussive  
Nasal Decongestant**

Sugar Free • Alcohol Free

Each dropperful (1.0 mL) contains:  
Chlorpheniramine Maleate . . . . . 1.0 mg  
Dextromethorphan HBr . . . . . 3.0 mg  
Phenylephrine HCl . . . . . 1.5 mg

1 fl oz (30 mL)

Lift Here for Drug Facts



N 3

## Drug Facts

### Active ingredients (in each 1 mL dropperful)

Chlorpheniramine Maleate 1.0 mg ..... Antihistamine  
Dextromethorphan Hydrobromide 3.0 mg ..... Antitussive (cough suppressant)  
Phenylephrine Hydrochloride 1.5 mg ..... Nasal Decongestant

### Purpose

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- nasal congestion
- swelling of nasal passages
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

Do not exceed recommended dosage. ▶

## Drug Facts (continued)

### Do not use this product

- if your child is now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if your child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- glaucoma
- a breathing problem such as chronic bronchitis
- a cough that lasts or is chronic such as occurs with asthma
- a cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking sedatives or tranquilizers. ▶

## Drug Facts (continued)

### When using this product

- may cause marked drowsiness
- sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.

### Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Lot: #Pxxxxx Exp. Date: xx/xxxx

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Supplied in a tight, light-resistant container with a child-resistant cap.

Manufactured for:  
**Lasar Pharmaceuticals, LLC**  
Greenville, SC 29615

Rev. 04/10

NDC 16477-811-30

# Donatussin DM DROPS

**Antihistamine  
Antitussive  
Nasal Decongestant**

Sugar Free • Alcohol Free

Each dropperful (1.0 mL) contains:

Chlorpheniramine Maleate . . . . . 1.0 mg  
Dextromethorphan HBr . . . . . 3.0 mg  
Phenylephrine HCl . . . . . 1.5 mg

1 fl oz (30 mL)



N 3

