

Donatussin Syrup

NDC 16477-186-01

Antihistamine * Cough Suppressant * Nasal Decongestant

Drug Facts

Active ingredients

(in each 5 mL teaspoonful)

Chlorpheniramine Maleate 4 mg

Dextromethorphan HBr 20 mg

Phenylephrine Hydrochloride 10 mg

473 mL

Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose and sneezing, itching of the nose or throat, and itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

_____ a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

a cough that occurs with too much phlegm (mucus)

heart disease

high blood pressure

thyroid disease

diabetes

trouble urinating due to an enlarged prostate gland

When using this product

- excitability may occur, especially in children.
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

_____Directions

Do not exceed 6 doses in a 24 hour period.

Adults and children 12 years of age and over: 1 teaspoonfuls (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours

Children 6 to under 12 years of age: ½ teaspoonful _____(2.5 mL) every _____4 hours, not to exceed 3 teaspoonfuls in _____24 hours

Children under 6 years of age: Consult a doctor

Other information

Store at 59° - 86°F (15° - 30°C)

Inactive ingredients

Berry Vanilla Flavor, Citric Acid, FD&C Red #40, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

Questions? Comments?

Call 1-864-286-8229



Manufactured for:

Laser Pharmaceuticals, LLC

Greenville, SC 29615

