

For full prescribing information, see the product foldout.

Store at 59° - 86°F (15° - 30°C)

Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing.

**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**

Dispense in a tight, light-resistant container as defined in the USP/NF with a child-resistant closure.

The labeling for this product includes professional labeling which is not intended for use by the general public.

**WARNING: KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN. IN CASE OF OVERDOSE, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.**

NDC: 16477-185-01

# Donatussin SYRUP

**New Improved  
Formula**

Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615

Patent Pending

Rev. 08/2011



**Antitussive  
Expectorant  
Nasal Decongestant**

Each 5 mL (1 teaspoonful) contains:

**Chlophedianol HCl . . . . . 12.5 mg**  
**Guaifenesin . . . . . 120 mg**  
**Phenylephrine HCl . . . . . 5 mg**



Berry Vanilla Flavor

**16 fl oz (473 mL)**

## **Drug Facts** (continued)

### **Other information**

Store at 59° - 86°F (15° - 30°C)

### **Inactive ingredients**

Berry Vanilla Flavor, Bitter Mask, Citric Acid, FD&C Red #40, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol, Sucralose.

### **Questions? Comments?**

Call 1-864-286-8229

Manufactured for:  
Laser Pharmaceuticals, LLC  
Greenville, SC 29615

# Drug Facts

Lift Here

## Active ingredients Purpose (in each 5 mL teaspoonful)

Chlophedianol  
Hydrochloride 12.5 mg ..... Antitussive  
Guaifenesin 120 mg ..... Expectorant  
Phenylephrine  
Hydrochloride 5 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- nasal congestion
- reduces swelling of nasal passages

## Warnings

**Do not exceed recommended dosage.**

### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- a cough that occurs with too much phlegm (mucus) ■ heart disease
- high blood pressure ■ thyroid disease
- diabetes

# Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

**Do not exceed 4 doses in a 24 hour period.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 6 to 8 hours, not to exceed 8 teaspoonfuls in 24 hours
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 6 to 8 hours, not to exceed 4 teaspoonfuls in 24 hours
Children 2 to under 6 years of age:	1/2 teaspoonful (2.5 mL) every 6 to 8 hours, not to exceed 2 teaspoonfuls in 24 hours
Children under 2 years of age:	Consult a doctor

4.125"