

NDC: 16477-562-01

# NEO DM Syrup

Tamper evident by foil seal under cap. Do not use if seal is broken or missing.

Dispense in a tight, light-resistant container with a child-resistant cap.



Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Greenville, SC 29615

Manufactured by:  
**Great Southern Laboratories**  
Houston, TX 77099

Rev. 10/09

**Antitussive  
Nasal Decongestant  
Antihistamine  
Sugar Free / Alcohol Free**

Each 5 mL (1 teaspoonful) contains:  
**Dextromethorphan HBr** ..... 30 mg  
**Pseudoephedrine HCl** ..... 50 mg  
**Brompheniramine Maleate** ..... 3 mg



16 fl oz (473 mL)

## Drug Facts

Lift Here

### Active ingredients Purpose (in each 5 mL teaspoonful)

Dextromethorphan	
Hydrobromide 30 mg	Antitussive
Pseudoephedrine	
Hydrochloride 50 mg	Nasal Decongestant
Brompheniramine	
Maleate 3 mg	Antihistamine

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- nasal congestion
- swelling of nasal passages
- runny nose
- itchy, watery eyes
- sneezing

### Warnings

**Do not exceed recommended dosage.**

#### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

## Drug Facts (continued)

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.**

#### When using this product

- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.  
Keep out of the reach of children.**

### Directions

**Do not exceed 4 doses in a 24 hour period.**

Adults and children 12 years of age and over:	1 teaspoonful every 6 hours.
Children 6 to under 12 years of age:	1/2 teaspoonful every 6 hours.
Children under 6 years of age:	Consult a physician.

## Drug Facts (continued)

### Other information

Store at 59° - 86°F (15° - 30°C)

### Inactive ingredients

Berry Vanilla Flavor, Citric Acid, FD&C Blue #1, FD&C Red #40, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

### Questions? Comments?

Call 1-864-286-8229