

LASER

NDC 16477-306-01

Respire[®]-30

Immediate-Release Capsules

Guaifenesin 150 mg
Pseudoephedrine Hydrochloride 30 mg

NASAL DECONGESTANT EXPECTORANT

Clears Nasal/Sinus Congestion
Thins and Loosens Mucus

100 CAPSULES

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL
OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

Drug Facts

Active Ingredients

(in each immediate-release capsule)

Guaifenesin 150 mg.....Expectorant
Pseudoephedrine HCl 30 mg.....Nasal decongestant

Uses

■ relieves nasal and sinus congestion and thin bronchial secretions to
improve breathing
■ relieves chest pain associated with coughing
■ relieves the common cold
■ relieves upper respiratory allergies
■ relieves sinusitis and/or sinus drainage
■ temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric
or emotional conditions, or Parkinson's disease), or for 2 weeks
after stopping the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor or pharmacist
before taking this product.

Manufactured by: Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale, NY 11735 308T 0309
Manufactured for: Laser Pharmaceuticals, LLC
Greenville, SC 29615



Lot#:

Exp. Date:

PEEL HERE
FOR MORE
DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if you have

■ heart disease ■ high blood pressure ■ thyroid disease

■ trouble urinating due to enlarged prostate gland

■ persistent or chronic cough such as occurs with smoking,
asthma, chronic bronchitis, or emphysema

■ cough accompanied by too much phlegm (mucus)

When using this product: ■ do not use more than directed

Stop use and ask a doctor if

■ you get nervous, dizzy or sleepless

■ symptoms do not get better within 7 days, come back or

occur with fever, rash or persistent headache. These could
be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional
before use.

Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right away.

Directions

■ Take with a full glass of water

■ This product can be administered without regard for timing
of meals

adults and children

1 or 2 capsules every

12 years of age and older 4-6 hours; not more than

8 capsules in 24 hours

children 6 to under

1 capsule every 4-6 hours;

12 years of age not more than 4 capsules in

24 hours

children under 6 years

do not use

of age

Other Information

■ Store at 25°C (77° excursions permitted between 15°-30°C
(59°-86°F)

■ use by expiration date on package

Inactive ingredients corn starch, D&C yellow #10,

FD&C blue #1, FD&C red #40, gelatin, iron oxide black,

povidone, silica, sugar, talc, titanium dioxide

Questions? Call 1-864-286-8229, 24 hours a day

STOP PEELING

ADHESIVE AREA