

**RONDEC-D- chlophedianol hydrochloride, pseudoephedrine hydrochloride liquid**  
**Laser Pharmaceuticals, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Rondec-D**

**Active ingredients**

**(in each 5mL teaspoonful)**

Chlophedianol Hydrochloride 12.5 mg

Pseudoephedrine Hydrochloride 30 mg

**Purpose**

Cough Suppressant

Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a physician before use if you have**

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**Stop use and ask a physician if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of the reach of children.**

**In case of accidental overdose, seek professional help or contact a Poison Control Center**

**immediately.**

***Directions***

**Do not exceed recommended dosage.**

---

Adults and children 12 years of age and over:	2 teaspoonfuls every 6 to 8 hours, not to exceed 8 teaspoonfuls in a 24 hour period
Children 6 to under 12 years of age:	1 teaspoonful every 6 to 8 hours, not to exceed 4 teaspoonfuls in a 24 hour period
Children under 6 years of age:	Consult a physician

---

***Other information***

Store at 59°-86°F (15°-30°C)

***Inactive ingredients***

Citric Acid, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

***Questions? Comments?***

Call 1-844-302-5227

**LASER**

Manufactured for:

Laser Pharmaceuticals, LLC

Nine North Drive, Suite 400

Alpharetta, GA 30004

1-844-302-5227

**Rondec-D 16 fl oz (473 mL) Bottle Label**

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Supplied in a tight, light-resistant container with a child-resistant cap.

This package not intended for dispensing to the patient.

NDC: 16477-101-16

# Rondec-D

## LASER

Manufactured for:  
**Laser Pharmaceuticals, LLC**  
Nine North Drive, Suite 400  
Alpharetta, GA 30004

US Patent # 9,050,289

Rev. 10/17



## Cough Suppressant/ Nasal Decongestant

Each teaspoonful (5 mL) for oral administration contains:

Chlophedianol HCl ..... 12.5 mg  
Pseudoephedrine HCl ..... 30 mg

Sugar Free/Alcohol Free

Dye Free/Gluten Free

## LASER

16 fl oz (473 mL)

### Drug Facts Lift Here

#### Active ingredients Purpose (in each 5 mL teaspoonful)

Chlophedianol  
Hydrochloride 12.5 mg .... Cough Suppressant  
Pseudoephedrine  
Hydrochloride 30 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

#### Warnings

**Do not exceed recommended dosage.**

#### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a physician or pharmacist before taking this product.

#### Ask a physician before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus) ■ heart disease
- high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland

**Drug Facts** (continued)**Stop use and ask a physician if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of the reach of children.**

**In case of an accidental overdose, seek professional help or contact a Poison Control Center immediately.**

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls every 6 to 8 hours, not to exceed 8 teaspoonfuls in a 24 hour period
Children 6 to under 12 years of age:	1 teaspoonful every 6 to 8 hours, not to exceed 4 teaspoonfuls in a 24 hour period
Children under 6 years of age:	Consult a physician

**Other information**

Store at 59° - 86° F (15° - 30° C)

**Inactive ingredients**

Citric Acid, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

**Drug Facts** (continued)**Questions? Comments?**

Call 1-844-302-5227

**LASER**

Manufactured for:  
Laser Pharmaceuticals, LLC  
Nine North Drive, Suite 400  
Alpharetta, GA 30004  
1-844-302-5227

**RONDEC-D**

chlorphedianol hydrochloride,pseudoephedrine hydrochloride liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:16477-101
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL
<b>CHLOPHEDIANOL HYDROCHLORIDE</b> (UNII: 69QQ58998Y) (CHLOPHEDIANOL - UNII:42C50P12AP)	CHLOPHEDIANOL HYDROCHLORIDE	12.5 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16477-101-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2017	
2	NDC:16477-101-15	15 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/21/2017	

**Labeler** - Laser Pharmaceuticals, LLC (614417132)

Revised: 11/2017

Laser Pharmaceuticals, LLC